

- [10] MARKAR S R, KARTHIKESALINGAM A, LOW D E. Outcomes assessment of the surgical management of esophageal cancer in younger and older patients [J]. Ann Thorac Surg, 2012, 94:1652–1660.
- [11] SCHELLERER V S, MERKEL S, SCHUMANN S C, et al. Despite aggressive histopathology survival is not impaired in young patients with colorectal cancer: CRC in patients under 50 years of age [J]. Int J Colorectal Dis, 2012, 27 (1):71–79.
- [12] LIM W H, DOGRA G K, MCDONALD S P, et al. Compared with younger peritoneal dialysis patients, elderly patients have similar peritonitis-free survival and lower risk of technique failure, but higher risk of peritonitis-related mortality [J]. Perit Dial Int, 2011, 31(6):663–71.
- [13] KEE Y K, PARK J T, YOON C Y, et al. Characteristics and clinical outcomes of end-stage renal disease patients on peritoneal dialysis for over 15 Years: a single-center experience [J]. Perit Dial Int, 2017, 37(5):535–541.
- [14] RUFINO J M, GARCIA C, VEGA N, et al. Current peritoneal dialysis compared with haemodialysis: medium-term survival analysis of incident dialysis patients in the Canary Islands in recent years [J]. Nefrologia, 2011, 31 (2):174–184.
- [15] PINES J M. Timing of antibiotics for acute, severe infections [J]. Emerg Med Clin North Am, 2008, 26 (2):245–257.
- [16] LEE C C, LEE C H, HONG M Y, et al. Timing of appropriate empirical antimicrobial administration and outcome of adults with community-onset bacteremia [J]. Crit Care, 2017, 21(1):119–127.
- [17] RYOO S M, KIM W Y, COHN C H, et al. Prognostic value of timing of antibiotic administration in patients with septic shock treated with early quantitative resuscitation [J]. Am J Med Sci, 2015, 349(4):328–333.
- [18] MUSZYNSKI J A, KNATZ N L, SARGEL CL, et al. Timing of correct parenteral antibiotic initiation and outcomes from severe bacterial community-acquired pneumonia in children [J]. Pediatr Infect Dis J, 2011, 30(4):295–301.
- [19] SARAH A S, RYAN W M, JASON R, et al. The impact of timing of antibiotics on outcomes in severe sepsis and septic shock: a systematic review and meta-analysis [J]. Crit Care Med, 2015, 43(9):1907–1915.
- [20] YE H, ZHOU Q, FAN L, et al. The impact of peritoneal dialysis-related peritonitis on mortality in peritoneal dialysis patients [J]. BMC Nephrol, 2017, 18(1):186.
- [21] 易春燕, 阳晓, 郭群英, 等. 长腹膜透析龄患者首次腹膜炎的临床特征分析 [J]. 中华肾脏病杂志, 2011, 27 (10):715–719.
- [22] ODUDU A, WILKIE M. Controversies in the management of infective complications of peritoneal dialysis [J]. Nephron Clin Pract, 2011, 118(3):C301–C308.
- [23] SZETO C C, CHOW K M. Gram-negative peritonitis—the Achilles heel of peritoneal dialysis? [J]. Perit Dial Int, 2007, 27(2):267–271.
- [24] BARRACLOUGH K, HAWLEY C M, MCDONALD S P, et al. Polymicrobial peritonitis in peritoneal dialysis patients in Australia: predictors, treatment, and outcomes [J]. Am J Kidney Dis, 2010, 55(1):121–131.
- [25] 刘小菁, 岳凌菊, 刘永泉, 等. 单中心腹膜透析相关性腹膜炎的临床分析 [J]. 中国血液净化, 2017, 16(4):251–254.

[本文编辑:刘晓华]

•编读往来•

## 通信作者的职责

通信作者是课题负责人,也是文章和研究的联系人,在论文投稿、修改直至发表的整个过程中,一切的联络工作均由其负责。通信作者掌握课题的经费,负责科研的设计以及文章的书写和质量把关,确保文章的真实性和科学性,是该论文的责任承担者。相对第一作者而言,通信作者具有更高的学术地位和专业水平,在该项科研中以第一作者的指导老师或重要辅导专家的身份为其提供帮助。对一篇科技论文来说,通信作者与第一作者所做的贡献是并列的,均为该文章最重要的作者。

[本刊编辑部]